

[Web](#) [Images](#) [Videos](#) [Maps](#) [News](#) [Shopping](#) [Gmail](#) [more](#) ▼

 Google scholar 

[Advanced Scholar Search](#)  
[Scholar Preferences](#)
**Scholar**   -   Results 21 - 30 of about 1,620. (0.11 sec)

[Helicobacter pylori uses motility for initial colonization and to attain robust infection- •nih.gov](#)

KM Ottemann, AC Lowenthal - **Infection** and immunity, 2002 - Am Soc Microbiol  
 ... storage, a thick 3- to 5-day growth of H. pylori from a Columbia blood agar plate was scraped into brucella broth with 10% FBS (BB10)-1% (wt/vol)  $\beta$ -cyclodextrin-25% glycerol-5 ... Antibody response to Campylobacter coli in children during intestinal **infection** and carriage. ...

[Cited by 78](#) - [Related articles](#) - [BL Direct](#) - [All 9 versions](#)
[Beta cyclodextrins enhance adenoviral-mediated gene delivery to the intestine- •umich.edu \(PDF\)](#)

MA Croyle, BJ Roessler, CP Hsu, R Sun, GL ... - Pharmaceutical research, 1998 - Springer  
 ... In this report, various cyclodextrin formulations were tested for their ability to enhance adeno ... replication-deficient adenovirus type 5 vectors encoded with either the E. coli beta-galactosidase ... was assessed by X-gal staining or visualization of fluorescence 48 hours after infection. ...

[Cited by 30](#) - [Related articles](#) - [BL Direct](#) - [All 6 versions](#)
[... of mast cell-mediated innate immunity during early response to bacterial infection](#)

R Malaviya, A Georges - Clinical Reviews in Allergy and Immunology, 2002 - Springer  
 ... encapsulation. Further,  $\beta$ -methyl cyclodextrin, an inhibitor of caveolae, blocked the uptake of FimH expressing E. coli in MCs. These ... viable. Modulation of MC Responses to Bacterial Infection by Immunoregulatory Cytokines and Growth Factors ...

[Cited by 35](#) - [Related articles](#) - [BL Direct](#) - [All 4 versions](#)
[Methods and compositions for treating or preventing bacterial infection](#)

D Zhu, M Muramatsu, J Xie, C Ni, MW ... - US Patent App. 09/975,136, 2001 - Google Patents  
 ... 5. [0071] 10: Petersen et al., Role of strain type, AGS cells and fetal calf serum in Helicobacter pylori adhesion and invasion ... specific embodiment, a method for treating or preventing a disease or disorder caused by bacterial **infection**, eg, E. coli or H. pylori **infection**, is provided ...

[Cited by 1](#) - [Related articles](#) - [All 7 versions](#)
[Type 1 fimbriated Escherichia coli-mast cell interactions in cystitis- •jstor.org \(PDF\)](#)

SN Abraham, JS Shin, R Malaviya - The Journal of infectious diseases, 2001 - UChicago Press  
 ... comes from the finding that specific disrupters of caveolae (methyl -  $\beta$  - cyclodextrin) or usurpers ... not uptake of inert beads or antibody - opsonized FimH - minus E. coli by mast ... mast cells was diverted to intracellular bacteria-containing chambers following bacterial **infection** of mast ...

[Cited by 12](#) - [Related articles](#) - [BL Direct](#) - [All 5 versions](#)
[Maltose/maltodextrin system of Escherichia coli: transport, metabolism, and ...- •nih.gov](#)

W Boos, H Shuman - Microbiology and Molecular Biology Reviews, 1998 - mmlr.highwire.org  
 ... The observation that MalM is also present in Salmonella typhimurium may suggest that this ... Of the five known MalT-dependent operons in E. coli, two (malPQ and malZ ... This is particularly obvious

with **cyclodextrins** and the p-nitrophenyl derivatives of maltooligosaccharides that ...

[Cited by 299](#) - [Related articles](#) - [BL Direct](#) - [All 12 versions](#)

### Subject Index to Volume 33

U pertusa Kjellm - Preparative Biochemistry and Biotechnology, 2003 - informaworld.com  
... of Escherichia coli, 321-339 Cell separation, continuous, by density gradient centrifugation,  
87-100 Chimeric synthetic peptide, from the core p19 protein and the envelope gp46 glycoprotein,  
use in the immunodiagnosis of HTLV-II virus infection, 29-38 b-Cyclodextrin-modified ...

[CITATION] ... Factor Domains and Anchorage in Internalization of Dr-Fimbriated Escherichia coli  
..., P Goluszko, V Popov, J Singhal, T ... - Infection and ..., 2000 - Washington: American Society for ...

... of Helicobacter pylori plays a role during colonization in a mouse model of infection- [nih.gov](#)


NR Salama, G Otto, L Tompkins, S Falkow - Infection and immunity, 2001 - iai.highwire.org  
... Columbia agar base (Oxoid), 5% defibrinated horse blood (HemoStat Labs), 0.2% -cyclodextrin  
(Sigma), 10 ... strain 342s V, which contained the aphA3 gene from C. coli, conferring kanamycin ...  
in the mixed infection, we could distinguish mutant bacteria from wild-type bacteria on ...

[Cited by 122](#) - [Related articles](#) - [BL Direct](#) - [All 8 versions](#)

... and expression of HopZ, a protein involved in adhesion of Helicobacter pylori- [nih.gov](#) (PDF)

B Peck, M Ortkamp, KD Diehl, E Hundt, B ... - Nucleic acids ..., 1999 - Oxford Univ Press  
... it is important to examine this option with regard to controlling a H.pylori infection. ... Antibodies  
directed against E.coli components were removed according to the method of Gruber and ... were  
grown for 48 h in Müller-Hinton broth containing 0.1% [beta]-cyclodextrin, harvested by ...

[Cited by 63](#) - [Related articles](#) - [BL Direct](#) - [All 8 versions](#)



Result Page: [Previous](#) [1](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#) [9](#) [10](#) [11](#) [12](#) [Next](#)

[Go to Google Home](#) - [About Google](#) - [About Google Scholar](#)

©2009 Google